Saving Tiny Heinies



Cavilon: Apply every 3 days OR when the patient receives a bath:

Due: _____

Last Applied: _____

On admission apply Cavilon No String Barrier. Reapply every 3 days and/or when baby receives a bath.
Let Cavilon dry for 60 seconds (make sure it is not tacky). This will create a barrier to protect the skin.
The skin may still look red under the Cavilon while it is healing, and the barrier remains in place.

If bleeding occurs or the rash is getting worse, please take a picture, put into medical record, and start using Desitin and contact Wound Care

Cleanse with sensitive wipes and apply Cavilon No Sting days	Barrier q 3
 For redness, non-yeast rash and intact skin, cleanse with wipes and apply Cavilon No Sting Barrier q 3 days. 	sensitive
 For <u>bleeding</u> and redness, cleanse with water wipes and cream with every diaper change. Please upload a pictu medical record before applying Desitin and contact W 	re into
 For erythematous, denuded (moist), non-yeast rash, clear water wipes, apply light dusting of stomahesive, spray with barrier spray to "crust" powder and apply Desitin into dia diaper change 	th cavilon
 For erythematous, with yeast rash (bumps), intact skin, of water wipes and apply Cavilon No Sting Barrier q 3 days 	eleanse with
 For erythematous, denuded (moist), with yeast (bumps) with water wipes and apply Cavilon No Sting Barrier q 3 c 	· · · · · · · · · · · · · · · · · · ·
If consult needed, any questions or elevations please real Wound Care. You can call Alyssa Railsback (M-F 8-5) at 2 send an email after 5p on weekends. I will continue to rounit during the week.	02-6253,

Saving Tiny Heinies

